arXiv:1212.3061v2 [quant-ph] 21 Dec 2012

Direct Detection of Classically Imperceptible Dark Matter through Quantum Decoherence

C. Jess Riedel
IBM Watson Research Center, Yorktown Heights, NY, USA

(Dated: December 24, 2012)

Astrophysical and cosmological observations suggest that the majority of mass in the universe is made up of invisible dark matter. Although various pieces of indirect evidence about its nature have been collected, the direct detection of dark matter has eluded experimental searches despite extensive effort. If the mass of dark matter is below 1 MeV, it is essentially imperceptible to conventional detection methods because negligible energy is transferred to nuclei during collisions. Here I propose directly detecting dark matter through the quantum decoherence it causes rather than its classical effects such as recoil or ionization. I show that quantum spatial superpositions are sensitive to low-mass dark matter which is inaccessible to classical techniques. More generally, this method enables the detection of hypothetical new phenomena which have no classical influence on normal matter whatsoever. This provides new independent motivation for the experimental pursuit of macroscopic superpositions, which have recently seen rapid progress.

It has been almost 80 years since dark matter (DM) was first proposed to explain the observed orbital velocities within galaxies and galaxy clusters. Much additional evidence for its existence has accumulated in the subsequent decades, but it has always been indirect and essentially gravitational. Existing experiments, observations, and theoretical preferences form a complicated thicket of conditional restrictions on any potential theory of DM [1, 2], but model-independent constraints are still rare and very valuable. Most desirable would be the *direct* detection of DM, that is, local experiments here on Earth that observe the interaction of the DM particle with the well-known protons, neutrons and electrons that compose the normal matter we see around us.

Based just on the movement of luminous matter in the galaxy and the virial theorem, we infer that DM forms a halo which has a density of roughly $\rho \sim 0.4$ GeV/cm^3 [3] in the neighborhood of the solar system. The DM particles should follow a non-relativistic Maxwellian velocity distribution centered around $v_0 \sim 230$ km/s in the galactic rest frame with a cutoff at the galactic escape velocity $v_\mathrm{e} \sim 600$ km/s [4]. Given this, any hypothetical mass m_DM for the DM particle then fixes its rough interaction rate with a particle of normal matter (either an electron or a nucleon) as a function of a scattering cross-section σ: $\Gamma_0 \sim v_0 \sigma \rho / m_\mathrm{DM}$. Typically, this is the total cross-section for spin-independent elastic scattering with nucleons.

Conventional DM direct detection experiments consist essentially of a large container of normal matter (e.g. liquid xenon) which is carefully watched for the tiny effects of an elastic collision with the DM particle such as recoil, vibration, heating, or ionization. These techniques rely on there being sufficient energy transfer from the DM to the target that the state of the target is substantially changed.

For a collision with a target particle of mass M, the energy deposited is of order $m_\mathrm{DM}^2 v_0^2 / M$. Most often the target is an atomic nucleus, for which direct detection experiments are sensitive down to a few keV of energy. This corresponds to a sensitivity to DM masses greater than a few GeV and lines up well with the Lee-Weinberg bound, which constrains the most popular form of weakly interacting massive particle (WIMP) dark matter to a mass of at least 2 GeV.

However, the most natural WIMP models have been challenged by galactic N-body simulations and negative direct experimental searches. It is prudent to allow for more general possibilities, and there are many proposals for sub-GeV mass such WIMPless [5], "MeV" [6], bosonic super-WIMP [7], or asymmetric [8] dark matter.

Calorimetry experiments which look for bulk heating rather than individual collisions have been able to explore down to 10 MeV in a modest cross-section range [9]. Scattering off of electrons—if it occurs—can probe masses as low as 1 MeV [10, 11] because the lighter electrons absorb more energy and also have a lower detectable energy threshold than nuclei. But conventional direct detection techniques probably cannot do any better than this. A 1 MeV DM candidate will deposit about an eV when colliding with an electron and less than 10^{-3} eV when colliding with a nucleus. For a keV candidate, the values are 10^{-6} and 10^{-9} eV, respectively. Classicaly, sub-MeV dark matter would be ghostly. Its dynamics could be strongly influenced by visible matter, but it would leave little trace.

In this article, I propose searching for low-mass DM by observing the quantum decoherence [12, 13] it causes rather than its direct influence on normal matter.



FIG. 1. **Decoherence detection with a Mach-Zehnder interferometer.** System $\mathcal{N}$ is placed in a coherent superposition of spatially displaced wavepackets which each travel a separate path and then are recombined. In the absence of system $\mathcal{D}$, the interferometer is tuned so that $\mathcal{N}$ will be detected at the bright port with near unit probability, and at the dim port with near vanishing probability. However, if system $\mathcal{D}$ scatters off $\mathcal{N}$, these two paths can decohere and $\mathcal{N}$ will be detected at the dim port 50% of the time.

DECOHERENCE WITHOUT CLASSICAL INFLUENCE

As an alternative to conventional direct detection methods, consider an atom interferometer which takes advantage of the de-Broglie-wave nature of matter. For now, let us specialize to a toy Mach-Zehnder interferometer, figure 1, although the ideas will apply generally. An atom $\mathcal{N}$ is prepared in a coherent superposition $|N_L\rangle + |N_R\rangle$ of two wavepackets, one taking the left path and one taking the right path, with something functioning as a beam splitter. These wavepackets are allowed to propagate over some length, and then they are recombined with a second splitter. Assuming the spread of the wavepackets is negligible, we can align the second splitter so that the atom ends up at one "bright" sensor with near unit probability, and at the other "dim" sensor with near vanishing probability.

The sensors effectively measure $\mathcal{N}$ in the basis $\{|N_\pm\rangle = |N_L\rangle \pm |N_R\rangle\}$, with $|N_+\rangle$ corresponding to the bright sensor and $|N_-\rangle$ to the dim sensor. Now we allow for the possibility of some hypothetical particle $\mathcal{D}$, which we call dark matter, which might be passing through the interferometer. Let the state $|D_\varnothing\rangle$ represent the absence of DM and take the evolution to be trivial when it is not present:

$$\Big[|N_L\rangle + |N_R\rangle\Big]|D_\varnothing\rangle \to \Big[|N_L\rangle + |N_R\rangle\Big]|D_\varnothing\rangle. \qquad (1)$$

Measuring in the basis $\{|N_\pm\rangle\}$ gives outcome $|N_+\rangle$ with certainty, as expected. But suppose the DM particle approaches in state $|D_{\rm in}\rangle$ and decoheres the superposition by scattering off the atom,

$$\Big[|N_L\rangle + |N_R\rangle\Big]|D_{\rm in}\rangle \to |N_L\rangle|D^{(L)}_{\rm out}\rangle + |N_R\rangle|D^{(R)}_{\rm out}\rangle, \qquad (2)$$

into the conditional states $|D^{(L)}_{\rm out}\rangle$ and $|D^{(R)}_{\rm out}\rangle$ with $\langle D^{(L)}_{\rm out}|D^{(R)}_{\rm out}\rangle \approx 0$, thereby recording which-path information. For low $m_{\rm DM}$, the phase-space localized wavepackets $|N_L\rangle$ and $|N_R\rangle$ of the atom are not significantly perturbed following the scattering event. But a measurement in the basis $\{|N_\pm\rangle\}$ now gives equal probability of either outcome. When the dim sensor clicks (which it will do half the time), this gives direct evidence of the dark matter $\mathcal{D}$ *even if it transfers negligible momentum to* $\mathcal{N}$.



FIG. 2. **Decoherence by dark matter with different de Broglie wavelengths.** An atom $\mathcal{N}$ in a superposition of spatial extent Δx is decohered by a DM particle $\mathcal{D}$ of wavelength λ. (a) In the short-wavelength limit $\lambda \ll \Delta x$, a single scattering event completely decoheres: $\gamma = \langle D^{(L)}_{\rm out}|D^{(R)}_{\rm out}\rangle \approx 0$. (b) For longer wavelengths, it takes many scattering events to decohere: $|\langle D^{(L)}_{\rm out}|D^{(R)}_{\rm out}\rangle| = 1 - \epsilon$, with ϵ small, but $|\gamma| = |\langle D^{(L)}_{\rm out}|D^{(R)}_{\rm out}\rangle|^N \approx e^{-N\epsilon} \approx 0$ for sufficiently large N.

This basic idea is suggested by the decoherence experiments of Hornberger et al. [14] and others. In the Hornberger et al. experiment, coherent spatial superpositions of C_{70} fullerenes were demonstrated by passing them through several gratings and recording the interference pattern. (This is a multi-slit near-field experiment rather than a Mach-Zehnder interferometer.) The interference region was filled with a gas of molecules much smaller than the fullerenes, such that collisions recorded which-path information in the gas but only mildly deflected the fullerenes. The pressure was adjustable so that the presence of the gas—and moreover its density—could be inferred from the suppression of the interference fringes. In that experiment, however, the deflection of the fullerenes was not negligible; the increasing pressure exponentially suppressed the count rate in addition to the fringe visibility.

COLLISIONAL DECOHERENCE BY DARK MATTER

The ability of a single DM particle in state $|D_{\rm in}\rangle$ to decohere an atom through elastic scattering is determined by the overlap $\langle D^{(L)}_{\rm out}|D^{(R)}_{\rm out}\rangle$ of the conditional dark-matter post-scattering states. This overlap, in turn, is strongly affected by the typical de Broglie wavelength $\lambda_0 = 2\pi\hbar/v_0 m_{\mathrm{DM}}$ of the DM particle compared to the spatial separation $\Delta\vec{x}$ between the two wavepackets. If $\Delta x \ll \lambda$, then $|\langle D_{\mathrm{out}}^{(L)}|D_{\mathrm{out}}^{(R)}\rangle|$ is only slightly away from unity. The DM cannot easily "see" the superposition, and it takes many scattering events to decohere the normal matter [15]. On the other hand, if $\Delta x \gg \lambda$, then $\langle D_{\mathrm{out}}^{(L)}|D_{\mathrm{out}}^{R)}\rangle$ nearly vanishes. The out states are distinct and a single DM collision suffices to decohere. See figure 2.

To calculate the state of the atom for the general case, we must sum the effect of all the DM particles passing by. So long as the DM is not degenerate, it can be treated as a fixed number of identically distributed but distinguishable particles. The state of the atom $\mathcal{N}$ in the $\{|N_L\rangle, |N_R\rangle\}$ basis after a time T is

$$\rho_{\mathcal{N}} = \frac{1}{2}\begin{pmatrix} 1 & \gamma \\ \gamma^* & 1 \end{pmatrix} \tag{3}$$

where $\gamma = \exp[-\int_0^T \mathrm{d}t\, F(\Delta\vec{x})]$ is the decoherence factor and $F(\Delta\vec{x})$ is given by [13, 16]

$$F(\Delta\vec{x}) = \int \mathrm{d}\vec{q}\, n(\vec{q}) \frac{q}{m_{\mathrm{DM}}} \int \mathrm{d}\hat{r} \times \left\{1 - \exp[i(\vec{q} - q\hat{r}) \cdot \Delta\vec{x}]\right\} |f(\vec{q}, q\hat{r})|^2 . \tag{4}$$

Here, $n(\vec{q})$ is the distribution function (phase space number density) of DM and $|f(\vec{q}_{\mathrm{in}}, \vec{q}_{\mathrm{out}})|^2 = \mathrm{d}\sigma/\mathrm{d}\Omega$ is the differential cross-section.

The condition for effective decoherence is that $\gamma \ll 1$, that is $\langle \mathrm{Re}\, F(\Delta x)\rangle \gtrsim 1/T$. This diagonalizes the density matrix and drives the probabilities for activating the sensors at either arm of the interferometer both to 1/2. For general q, the angular integrals over $\hat{q}$ and $\hat{r}$ in (4) can only be done by assuming a form for the differential cross-section. For the purpose of calculating dark matter sensitivity, I will assume hard-sphere scattering, i.e. $\mathrm{d}\sigma/\mathrm{d}\Omega$ constant. So long as the cross-section does not vary too quickly with momentum q, the bounds will not be very sensitive to this simplification.

MASSIVE SUPERPOSITIONS

The toy detector in figure 1 only works if the flux of DM is high enough such that at least one DM particle will usually scatter off of an atom wavepacket while it is in the interferometer. Because of the rarity of collisions, interferometry with single atoms has little hope of being sensitive to DM. There are at least two ways to increase the likelihood of a scattering event, and hence increase the possibility of detection. First, the time over which the superposition is maintained can be increased, by lengthening the interferometer arms or slowing down the atom. For an uncorrelated flux of DM, the expected number of scattering events should be linear in time. Second, and more powerfully, one can superpose ever larger clusters of atoms. That is, construct a *matter* interferometer with targets $\mathcal{N}$ which are as large as possible. As the number of nucleons composing $\mathcal{N}$ increases, each contributes an independent decoherence factor. This multiplies the effective decoherence rate $\mathrm{Re}\, F(\Delta\vec{x})$ by the total number of nucleons.



FIG. 3. **Coherent scattering over multiple nuclei.** (a) When the target object is composed of many atoms, short-wavelength DM can scatter off a single nucleus. The total cross-section is proportional to the mass of the target. (b) For sufficiently small m_{DM}, the DM wavelength is too long to resolve the individual nuclei. Instead, the DM scatters off the entire target coherently and the total cross-section is proportional to the target mass *squared*.

Furthermore, sub-GeV DM will scatter coherently off of multiple nucleons if its coupling to them is spin-independent, as is commonly assumed. This means that the total scattering rate is proportional to the square of the number of nucleons in the target when the target is smaller than the reduced de Broglie wavelength $\bar{\lambda} = \lambda/2\pi$ of the DM, i.e. the target fits inside the coherence volume $\bar{\lambda}^3$. See figure 3. (DM essentially always scatters coherently off of multiple nucleons in the same nucleus for $m_{\mathrm{DM}} < 1$ GeV.) When the wavelength is large enough to span multiple nuclei, $m_{\mathrm{DM}} \lesssim 1$ MeV, but not large enough to span the whole target, complicated interference effects (both constructive and destructive) arise which depend on the distribution of nuclei and the form of the angular cross-section. In this article, I take the coherence boost compared to incoherent scattering to be proportional to the number of nuclei which fit in the coherence volume.

To achieve interference of large objects with ever smaller de Broglie wavelengths, modern time-domain interferometers can require a time interval proportional to the size of the object superposed [17]. When combined with the coherence boost, the DM sensitivity scales like the *cube* of the mass of the object. Although this is partially a testament to the difficulty of superposing large objects, it also means that investing in larger masses yields big dividends. Happily, recent progress in the size of superposed objects in matter interferometry has been stunning, with clear fringe patterns produced when interfering molecules composed of up to 430 atoms and in excess of 6,000 amu [18]. Future prospects are even stronger [17, 19], and these have great potential for discovery.

Of course, there are many possible sources of decoherence; anomalous decoherence hardly implies the existence of new particles. Still, note that the inverse statement *is* true: the observation of interference effects (which establishes the existence of the superposition, and hence implies that all relevant sources of decoherence have been eliminated) implies that DM has *not* scattered (which sets robust upper bounds on the cross section σ). Furthermore, if anomalous decoherence is observed, one may be able to demonstrate that it is due to DM by observing the functional dependence of the interference fringe visibility on parameters which affect the decoherence rate. For instance, the spatial extent Δx of the superposition can be adjusted by separating the arms of an interferometer, while the exposure time T can be changed by varying the length of the arms or the speed of the target. Depending on the design, varying the isotopic composition might allow one to adjust the dark-matter cross-section of the nuclei without affecting other potential sources of decoherence. Finally, the DM flux itself might be manipulable, which is discussed below.

COHERENT EFFECTS AND MODULATION

Unlike earlier examples of collisional decoherence [13, 15, 16], the scattering DM environment is not distributed isotropically because of the significant speed with which the sun orbits the galactic center. The apparent DM "wind" drives the imaginary part of $F(\vec{\Delta x})$ to a non-zero value, a result of forward scattering [20].

$\operatorname{Im} F(\vec{\Delta x})$ changes the phase of the decoherence factor γ. When $\operatorname{Re} F(\vec{\Delta x})$ is large compared to T, this doesn't matter; γ vanishes regardless. However, when $\operatorname{Re} F(\vec{\Delta x})$ is small, $|\gamma| \approx 1$ and the density matrix is then given by $\rho \approx |\psi\rangle\langle\psi|$ for $|\psi\rangle = |N_L\rangle + \gamma|N_R\rangle$. The state has not been decohered and is still pure. Instead, the DM environment has acted unitarily on the normal matter by applying a position-dependent phase.

A position-dependent phase per unit time is simply a coherent *classical* force, and it is natural that this net force is only non-zero when the DM momentum distribution is anisotropic. Although the DM wind does not measurably displace the wavepackets $|N_L\rangle$ and $|N_R\rangle$, it is known that such forces can still be detected using matter interferometry. This is a quantum-enhanced measurement [21, 22] and, although it has not been used to detect new particles, it is the basis behind many existing weak-force experiments (e.g references [23, 24]). The details of the DM and the interferometer determine the ratio $\operatorname{Im} F / \operatorname{Re} F$, and this governs the transition from the well-known coherent case (which can be modeled as a unitary evolution) to the decoherent case introduced here (which cannot).

Since interferometers cannot measure a constant phase shift between their two arms, the force must vary to be observable. There are therefore two related motivations for modulating the influence of the DM: (1) if anomalous decoherence is detected, its functional dependence on parameters which control the hypothetical DM scattering gives evidence that the decoherence is in fact due to DM, and (2) if the DM wind applies only a coherent phase shift, rather than decoherently dephasing, then some variation in time is necessary to observe this shift at all. Several strategies are possible.

First, the incoming DM may be directly shielded from reaching the detector. This can be done using normal materials, such as lead or concrete, for most of the relevent parameter space. Alternatively, for low cross-sections, the detector could be operated underground.

Second, predictable natural variations in the apparent DM flux, such as those due to the Earth's orbital motion around the sun, may be exploited. This technique has an extensive history [25–27].

Third, outside of the short wavelength limit (i.e. when it takes multiple scattering events to fully decohere), the orientation of the superposition with respect to the direction of the DM wind gives an order-unity modulation of the effect. Decoherence is usually maximized when the wind is parallel to the separation vector $\vec{\Delta x}$. Alternatively, when the coherent effect dominates, the phase shift due to the wind flips sign as the orientation is rotated. See figure 4. This means interferometers are naturally *directional* DM detectors, which are known to be highly desirable [28] in part because they provide unmistakable evidence that a signal is of galactic origin. The Earth's daily rotation guarantees that this directional variation, combined with the effects of the Earth's DM wind shadow, will be visible even to fixed terrestrial experiments.

DARK MATTER SEARCH POTENTIAL

The search potential for low-mass DM through decoherence is depicted in figure 5. I will concentrate on the wide mass range 10 eV–100 MeV. Above this range, conventional direct detection techniques will be superior. Below this range, the occupation number of DM momentum modes in the Milky Way surpasses unity. (Fermionic DM would show signs of degeneracy, while bosonic DM would behave like a coherent wave.)

The only existing direct-detection bound on the spin-independent nucleon-scattering cross-section for $m_{\mathrm{DM}} < 1$ GeV comes from the X-ray Quantum Calorimetry experiment [36], as analyzed by Erickcek et al. [9]. There is also a constraint arising from the stability of the DM halo which encompasses the Milky Way, and the rate at which it heats interstellar hydrogen through collisions [29]. These robust exclusions are based only on the



FIG. 4. **Visibility of the dark matter wind.** (a) The Earth moves with $v_{\text{Earth}} \sim 230$ km/s in the Milky Way. (b) Matter passing through an interferometer is in a superposition of two wavepackets separated by $\vec{\Delta x}$. (c) The DM momentum distribution (red arrows) seen by the experiment is concentrated in a direction opposite $\vec{v}_{\text{Earth}}$. The sensitivity of the superposition to interactions with DM is determined by the angle χ between $\vec{v}_{\text{Earth}}$ and $\vec{\Delta x}$. χ can be adjusted by re-orienting $\vec{\Delta x}$, i.e. rotating the interferometer. (d) The real part of F controls the decoherence of the superposition. It is plotted for $d\sigma/d\Omega \propto 1$ (hard-sphere) scattering as a function of χ and $\zeta = m_{\text{DM}} v_0 \Delta x/\hbar = \Delta x/\lambda_0$. (It is normalized to its value at $\chi = 0$.) Values for $\chi > \pi/2$ are given by $\text{Re}\, F_{\pi-\chi} = \text{Re}\, F_\chi$. For DM wavelengths much shorter than the size of the superposition ($\zeta \to \infty$), there is no dependence on orientation because a single collision event causes complete decoherence. The insets compare χ dependence for $d\sigma/d\Omega \propto 1$ scattering (blue) with $d\sigma/d\Omega \propto 1 + \cos^2\theta$ (red) and $d\sigma/d\Omega \propto \cos^4\theta$ (green) for fixed ζ. (The vertical scale varies between insets.) This is how the decoherence strength would fluctuate as an interferometer is rotated with respect to the DM wind. (e) The imaginary part of F controls the coherent phase shift between the arms of the interferometer due to the weak force applied by the wind. Values for larger χ are given by $\text{Im}\, F_{\pi-\chi} = -\text{Im}\, F_\chi$. Although the phase shift has χ dependence for all ζ, decoherence always prevents the observation of this shift in the short-wavelength limit: $\text{Im}\, F_\chi/\text{Re}\, F_\chi \to 0$ as $\zeta \to \infty$.

present-day distribution of the DM which is necessary to explain observed galactic dynamics.

If one further assumes the simplest thermal freeze-out scenario for DM in the early universe, tighter upper bounds on σ for $m_{\text{DM}} > 1$ MeV have been derived from the cosmic microwave background and large-scale structure data [30]. This type of thermal relic DM also becomes too warm to explain small-scale structure data encoded in the Lyman-α forest when the mass falls below roughly a keV (e.g. [31]). Both of these restrictions might easily be violated (by dark-observable temperature ratios [5] or fully non-thermal scenarios), as might other published bounds which rest on significant new assumptions about the nature of DM (e.g. its self-annihilation or high-energy inelastic scattering to gamma rays [37]).

Direct-detection experiments on Earth will only be sensitive to DM if the scattering cross-section with nucleons is sufficiently low for DM to pass through the atmosphere and reach the experiment. This upper bounds the spin-independent cross-section visible to experiments on



FIG. 5. **The sensitivity of several existing and proposed superposition experiments to the spin-independent nucleon scattering cross-section of dark matter, compared with existing constraints.** (a) Gray shaded regions are robustly excluded by the X-ray Quantum Calorimetry experiment ("XQC" [9]) and galactic heating and halo stability arguments ("Heating+Halo" [29]). Hatched regions are incompatible with thermal DM models due to observations of the cosmic microwave background with large scale structure data ("CMB+LSS" [30]) and the Lyman-α forest ("Ly-α" [31]). Solid colored lines bound regions where DM would cause decoherence in three proposed experiments: a satellite-based atom interferometer ("AGIS" [24]), optically-trapped silicon nanospheres ("Nanosphere" [19]), and the OTIMA interferometer with clusters of gold of mass 10^6 amu ("OTIMA-6" [17]). A successful AGIS satellite would set new exclusion limits on DM where its sensitivity dips below the halo heating/stability bound for $m_{\rm DM} \lesssim 1$ keV. On the other hand, the OTIMA and nanosphere experiments would be shielded from DM by the atmosphere if operated at sea level, so exclusion regions illustrate the sensitivity at an altitude of 200 km. The darker regions bordered by colored dashed lines indicates where the coherent phase shift due to the DM wind could be observed without being overwhelmed by decoherence. (b) On top of the existing exclusions (now black dotted lines), the colored lines give the lower limits on the sensitivities of existing interferometers with helium atoms ("He" [32]), cold neutrons ("n^0" [33]), fullerenes ("C$_{70}$" [34]), and the large organic molecule C$_{60}$[C$_{12}$F$_{25}$]$_{10}$ ("PFNS10" [18]). Also shown are sensitivities for the AGIS satellite, the OTIMA interferometer with three choices of gold cluster mass (10^4, 10^6, and 10^8 amu, although the last is not feasible for an Earth-bound experiment), and the nanosphere experiment. The border is defined by an e-fold suppression of the interference fringes: $|\gamma| = 1/e$. Sensitivity increases dramatically for larger masses. When operated within the Earth's atmosphere, there is a potential to detect DM only where the sensitivity dips below the dashed-dotted line corresponding to the degree of shielding at the relevant altitudes. None of the existing experiments do so. For reverence, strongly interacting massive particle ("SIMP" [35]) models lie in the black band.

the Earth's surface at about 10^{-27}cm^2. This can be circumvented by placing the experiment on a high-altitude balloon (~30 km altitude; ~10^{-25}cm^2), a sounding rocket (~200 km altitude; ~10^{-18}cm^2), or on a satellite. I do not know if anyone has studied the possibility of producing macroscopic quantum superpositions on a balloon or rocket, but spaceborne experiments are both feasible and compelling for independent reasons [17, 38, 39].

Figure 5 shows the potential reach of several existing matter interferometers [18, 32–34] in the absence of atmospheric shielding. The separation vector $\vec{\Delta x}$ is assumed to point into the DM wind. The effects of rotating $\vec{\Delta x}$ with respect to the wind are order unity and are depicted in figure 4. Modern experiments often use multiple gratings with many slits to overcome difficulties with beam coherence and tiny de Broglie wavelengths [40, 41], so the matter is not described by a simple superposition of two spatially separated wavepackets. But the interferometers still require good coherence over distances which span multiple slits, so it is reasonable to estimate their sensitivity by taking Δx to be the period of the relevant grating. (Only the results for small $m_{\rm DM}$ will depend on the choice of Δx; for larger masses, which are in the short wavelength limit, any scattering event results in complete decoherence independent of the spatial separation.)

To demonstrate the potential of future experiments to detect DM through decoherence, I consider three proposals currently being developed: an optical time-domain ionizing matter-wave (OTIMA) interferometer which will interfere clusters of atoms larger than 10^6 amu [17], an optically-trapped 40 nm silicon nanosphere with mass $\sim 10^8$ amu [19], and a satellite-based Atomic Gravitational wave Interferometric Sensor (AGIS) [24] using an estimated exposure time of 100 s on rubidium-87.

The AGIS satellite would interfere atoms in the open vacuum of space, so it would be sensitive to DM scenarios for $m_{\rm DM} < 1$ keV which have never been excluded, as shown in figure 5. The OTIMA and nanosphere experiments would need to be raised at least partially out of the Earth's atmosphere to see dark matter, but experiments further into the future could rule out DM scenarios at ground level. For $m_{\rm DM} < 3$ keV, there is a significant region for which DM's unitary phase shift can be observed in the OTIMA and nanosphere experiments without being overwhelmed by decoherence.

It is likely [42] that the true DM velocity distribution has a thicker tail than the Maxwellian form assumed here. High momentum DM causes disproportionately more decoherence, so this should increase sensitivity further. Also note that if there were a mechanism which increased the local DM density in the vicinity of the Earth (e.g. reference [43]), this would improve of the sensitivity of decoherence detection and other near-Earth methods without changing the astrophysical limits which currently provide the best bounds.

DISCUSSION

It's worth stressing that detection through decoherence is not limited to interferometers. Macroscopic superpositions of mechanical oscillators [44, 45] are promising because of the sheer size of the masses under quantum control. These are very different than the traditional interferometers depicted in figure 5 because the spatial separation of the superposition is much smaller than the physical size of the relevant objects. Rough estimates suggest that some of these proposed devices would have even larger sensitivity to DM, although a detailed analysis has not been performed.

The scattering cross-section of DM with electrons, rather than nucleons, is also of interest [10, 11]. It could be probed with the matter interferometers discussed here and also possibly with superconducting qubits, in which two experimentally manipulable quantum states are composed of millions of entangled Cooper pairs. Scattering DM could record "which-momentum" information about the electrons in these macroscopic states, and so decohere them. On the other hand, Bose-Einstein condensates (BECs) are not naturally suited for detection by decoherence. Spatial interferometry has been done with BECs [46], but this is essentially *atom* interferometry; the atoms in a BEC are all in the same state, but they aren't entangled. Individual atoms in the BEC can be lost without destroying the coherence, so there is no boost in sensitivity like there is for superposing large clusters of atoms. (For the same reason, normal BECs do not avoid the shot noise limit when used to measures phase differences.) However, the creation of entanglement in BECs, such as NOON [47] or spin-squeezed [48] states, might be exploitable for detection by decoherence.

All of the experiments discussed in this article were performed or proposed for reasons that had nothing to do with discovering new particles; significant enhancement in sensitivity is likely for devices designed with it in mind. Beyond dark matter, one can reinterpret many experiments which establish certain quantum states as direct evidence against hypothetical weak phenomena that, if existent, would decohere those states. The toy Mach-Zehnder interferometer illustrates that the classical effects of such phenomena (e.g. momentum transfer) can be arbitrarily small while still causing very noticeable decoherence.

The essential difficulty in creating macroscopic superpositions is that the interaction of a single constituent particle is enough to decohere an arbitrarily large system, but this extreme sensitivity also gives them their detecting power. Insofar as stability in the presence of decoherence defines the classicality of a quantum state [12, 49], the best constraints on hypothetical weak phenomena will come from the most non-classical states. This gives new independent motivation for their experimental pursuit.

Acknowledgements

I thank Markus Arndt, Asimina Arvanitaki, Dirk Bouwmeester, Xiaoyong Chu, Savas Dimopoulos, Rouven Essig, Alexander Friedland, Jay Gambetta, Christian Hagmann, Steen Hannestad, Andrew Hime, Lorenzo Maccone, Gregory Mack, Jeremy Mardon, Benjamin Monreal, Kevin Moore, Harry Nelson, Bryon Neufeld, Shmuel Nussinov, Keith Rielage, Daniel Sank, Robert Scherrer, Alexia Schulz, Mark Srednicki, Paul Steinhardt, Lev Vaidman, Neal Weiner, and Haibo Yu for useful discussion. I am especially grateful to Michael Graesser for teaching, Godfrey Miller for criticism, and Charlie Bennett, Jim Hartle, and Wojciech Zurek for making this work possible. This research was partially supported by the U.S. Department of Energy through the LANL/LDRD program, and by the John Templeton Foundation.

---

[1] Bartelmann, M. The dark universe. *Reviews of Modern Physics* **82**, 331 (2010).

[2] Beringer, J. *et al.* Review of particle physics. *Physical Review D* **86**, 010001 (2012).

[3] Catena, R. & Ullio, P. A novel determination of the local dark matter density. *Journal of Cosmology and Astroparticle Physics* **2010**, 004 (2010).

[4] Lewin, J. & Smith, P. Review of mathematics, numerical factors, and corrections for dark matter experiments based on elastic nuclear recoil. *Astroparticle Physics* **6**, 87 – 112 (1996).

[5] Feng, J., Tu, H. & Yu, H. Thermal relics in hidden sectors. *Journal of Cosmology and Astroparticle Physics* **2008**, 043 (2008).

[6] Hooper, D. & Zurek, K. M. Natural supersymmetric model with mev dark matter. *Phys. Rev. D* **77**, 087302 (2008).

[7] Pospelov, M., Ritz, A. & Voloshin, M. Bosonic superwimps as kev-scale dark matter. *Phys. Rev. D* **78**, 115012 (2008).

[8] Falkowski, A., Ruderman, J. & Volansky, T. Asymmetric dark matter from leptogenesis. *Journal of High Energy Physics* **2011**, 1–32 (2011).

[9] Erickcek, A. L., Steinhardt, P. J., McCammon, D. & McGuire, P. C. Constraints on the interactions between dark matter and baryons from the x-ray quantum calorimetry experiment. *Phys. Rev. D* **76**, 042007 (2007).

[10] Essig, R., Mardon, J. & Volansky, T. Direct detection of sub-gev dark matter. *Phys. Rev. D* **85**, 076007 (2012).

[11] Graham, P. W., Kaplan, D. E., Rajendran, S. & Walters, M. T. Semiconductor probes of light dark matter. *Physics of the Dark Universe* **1**, 32–49 (2012).

[12] Zurek, W. H. Decoherence, einselection, and the quantum origins of the classical. *Rev. Mod. Phys.* **75**, 715–775 (2003).

[13] Schlosshauer, M. *Decoherence and the Quantum-to-Classical Transition* (Springer-Verlag, Berlin, 2008).

[14] Hornberger, K. *et al.* Collisional decoherence observed in matter wave interferometry. *Phys. Rev. Lett.* **90**, 160401 (2003).

[15] Joos, E. & Zeh, H. D. The emergence of classical properties through interaction with the environment. *Zeitschrift für Physik B Condensed Matter* **59**, 223–243 (1985).

[16] Gallis, M. R. & Fleming, G. N. Environmental and spontaneous localization. *Phys. Rev. A* **42**, 38–48 (1990).

[17] Nimmrichter, S., Haslinger, P., Hornberger, K. & Arndt, M. Concept of an ionizing time-domain matter-wave interferometer. *New Journal of Physics* **13**, 075002 (2011).

[18] Gerlich, S. *et al.* Quantum interference of large organic molecules. *Nat Commun* **2** (2011).

[19] Romero-Isart, O. *et al.* Large quantum superpositions and interference of massive nanometer-sized objects. *Phys. Rev. Lett.* **107**, 020405 (2011).

[20] Schmiedmayer, J. *et al.* Index of refraction of various gases for sodium matter waves. *Phys. Rev. Lett.* **74**, 1043–1047 (1995).

[21] Braginsky, V. B., Khalili, F. Y. & Thorne, K. S. *Quantum Measurement* (Cambridge University Press, Cambridge, UK, 1992).

[22] Giovannetti, V., Lloyd, S. & Maccone, L. Quantum-enhanced measurements: Beating the standard quantum limit. *Science* **306**, 1330–6 (2004).

[23] Peters, A., Chung, K. & Chu, S. Measurement of gravitational acceleration by dropping atoms. *Nature* **400**, 849–852 (1999).

[24] Dimopoulos, S., Graham, P. W., Hogan, J. M., Kasevich, M. A. & Rajendran, S. Atomic gravitational wave interferometric sensor. *Phys. Rev. D* **78**, 122002 (2008).

[25] Michelson, A. A. & Morley, E. W. On the relative motion of the earth and the luminiferous ether. *American Journal of Science* **34**, 333–345 (1887).

[26] Aalseth, C. E. *et al.* Search for an annual modulation in a *p*-type point contact germanium dark matter detector. *Phys. Rev. Lett.* **107**, 141301 (2011).

[27] Bernabei, R. *et al.* Dark Matter search. *Nuovo Cimento Rivista Serie* **26**, 010000–74 (2003).

[28] Ahlen, S. *et al.* The case for a directional dark matter detector and the status of current experimental efforts. *International Journal of Modern Physics A* **25**, 1–51 (2010).

[29] Chivukula, R. S., Cohen, A. G., Dimopoulos, S. & Walker, T. P. Bounds on halo-particle interactions from interstellar calorimetry. *Phys. Rev. Lett.* **65**, 957–959 (1990).

[30] Chen, X., Hannestad, S. & Scherrer, R. J. Cosmic microwave background and large scale structure limits on the interaction between dark matter and baryons. *Phys. Rev. D* **65**, 123515 (2002).

[31] Narayanan, V., Spergel, D., Dave, R. & Ma, C.-P. Constrains on the mass of warm dark matter particles and the shape of the linear power spectrum from the lyα forest. *The Astrophysical Journal Letters* **543**, L103–L106 (2000).

[32] Carnal, O. & Mlynek, J. Young's double-slit experiment with atoms: A simple atom interferometer. *Phys. Rev. Lett.* **66**, 2689–2692 (1991).

[33] Zeilinger, A., Gähler, R., Shull, C. G., Treimer, W. & Mampe, W. Single- and double-slit diffraction of neutrons. *Rev. Mod. Phys.* **60**, 1067–1073 (1988).

[34] Brezger, B. *et al.* Matter-wave interferometer for large molecules. *Phys. Rev. Lett.* **88**, 100404 (2002).

[35] Wandelt, B. *et al.* Self-interacting dark matter. In *Sources and Detection of Dark Matter and Dark Energy in the Universe*, vol. 1, 263 (2001).

[36] McCammon, D. *et al.* A high spectral resolution observation of the soft x-ray diffuse background with thermal detectors. *The Astrophysical Journal* **576**, 188 (2002).

[37] Gregory D. Mack, A. M. Closing the window on strongly interacting dark matter (2012). arXiv:1211.1951.

[38] Kaltenbaek, R., Hechenblaikner, G., Kiesel, N., Johann, U. & Aspelmeyer, M. Macroscopic quantum resonators in space. In *Lasers and Electro-Optics Europe (CLEO EUROPE/EQEC), 2011 Conference on and 12th European Quantum Electronics Conference*, 1 (2011).

[39] Kaltenbaek, R. *et al.* Macroscopic quantum resonators (MAQRO). *Experimental Astronomy* 1–42 (2012).

[40] Cronin, A. D., Schmiedmayer, J. & Pritchard, D. E. Optics and interferometry with atoms and molecules. *Rev. Mod. Phys.* **81**, 1051–1129 (2009).

[41] Hornberger, K., Gerlich, S., Haslinger, P., Nimmrichter, S. & Arndt, M. *Colloquium*: Quantum interference of clusters and molecules. *Rev. Mod. Phys.* **84**, 157–173 (2012).

[42] Kuhlen, M. *et al.* Dark matter direct detection with non-maxwellian velocity structure. *Journal of Cosmology and*

Astroparticle Physics **2010**, 030 (2010).

[43] Adler, S. L. Can the flyby anomaly be attributed to earth-bound dark matter? *Phys. Rev. D* **79**, 023505 (2009).

[44] OConnell, A. *et al.* Quantum ground state and single-phonon control of a mechanical resonator. *Nature* **464**, 697–703 (2010).

[45] Romero-Isart, O., Clemente, L., Navau, C., Sanchez, A. & Cirac, J. I. Quantum magnetomechanics with levitating superconducting microspheres. *Phys. Rev. Lett.* **109**, 147205 (2012).

[46] Schumm, T. *et al.* Matter-wave interferometry in a double well on an atom chip. *Nat. Phys.* **1**, 57–62.

[47] Cable, H., Laloë, F. & Mullin, W. J. Formation of noon states from fock-state bose-einstein condensates. *Phys. Rev. A* **83**, 053626 (2011).

[48] Riedel, M. F. *et al.* Atom-chip-based generation of entanglement for quantum metrology. *Nature* **464**, 1170–3.

[49] Zurek, W. H. Pointer basis of quantum apparatus: Into what mixture does the wave packet collapse? *Phys. Rev. D* **24**, 1516–1525 (1981).